

The Casimir energy for two-dimensional deformed sphere

N. Shtykov

*Department of Physics , Tokyo Institute of Technology,
Oh-Okayama, Meguro-ku, Tokyo, 152, Japan[†]*

and

D. V. Vassilevich

*Department of Theoretical Physics, St. Petersburg University,
198904 St.Petersburg, Russia*

Abstract

We compute the Casimir energy for a free scalar field on the spaces $R^{m+1} \times \tilde{S}^2$ where $\tilde{S}^2$ is two-dimensional deformed two-sphere.

[†] On leave from: *Irkutsk university, Russia;*

Electronic mail: shtykov@phys.titech.ac.jp

The explanation of the Casimir effect lies in the fact, that quantum fields respond to the presence of external constraints. Meaningful constraints can be defined, for example, in terms of the boundary conditions which fix these fields. In a number of works, the Casimir energy has been calculated in d -dimensional spaces for free scalar, spinor and vector fields at a zero and non-zero temperature with the Dirichlet, von Neumann and Robin boundary conditions and the periodicity condition (compactification R^1 to S^1) [1]-[6]. (For review of several applications of the Casimir energy see ref. [7]).

The calculation of the one-loop effective potential (which is closely related to the Casimir energy) in product spaces $R^n \times Y$ where Y is a compact manifold was motivated by the Kaluza-Klein models and involves some mathematical problems related to regularization of the functional determinants , $\det \Delta$ with the eigenvalues of a second-order differential operator Δ [8]. The scalar functional determinants have been also determined in terms of ζ functions on sectors of the two-dimensional disk and spherical cap [9]. Recently, there has been some interest in investigating spaces with conical singularities. Physical applications for such spaces range from quantum cosmology [10] and cosmic strings [11] to finite temperature field theory [12]. Some mathematical aspects were examined in the papers [13]. Typically, two alternative cases are considered. In the first case, the conical singularities come from factorization of a sphere S^d with respect to a finite group Γ , which acts on S^d with fixed points. On this way, only discrete series of conical singularities can be studied. In the second case, a standard two-dimensional cone with varying angle deficit is introduced. However, in this approach curvature can be considered only as perturbation of the flat case. We propose to consider a two-dimensional sphere with two conical singularities with an arbitrary angle parameter. This can be treated as an imitation of the effects [12] for essentially curved space without exact relevance to any particular physics. In this sence our aim is two-fold. First, we study technical aspects of the problem and second, we explore qualitative behaviour of the effective potential which appears to be quite different from the non-singular case.

In this letter we compute the vacuum energy for free scalar fields with a background geometry of $R^{m+1} \times \tilde{S}^2$ where R^{m+1} , is the $m + 1$ -dimensional Euclidean space, $\tilde{S}^2$ is the two-dimensional compact manifold constructed from two-sphere S^2 by the deformation of its metric. We find the exact eigenvalues of the Laplace operator and compute the vacuum energy as a function of the deformation parameter by making use of the zeta-function technique. In odd-dimensional spaces $R^1 \times \tilde{S}^2$ and $R^3 \times \tilde{S}^2$, we obtain the finite vacuum energy, which changes its sign with

some value of the deformation parameter. In even dimensional spaces $R^2 \times \tilde{S}^2$ and $R^4 \times \tilde{S}^2$ the vacuum energy contains a divergent part and has the following form

$$E = \frac{1}{\epsilon} f_0(m, \rho) + f_1(m, \rho) + f_0 \log(r\mu) \quad (1)$$

where r is the radius of $\tilde{S}^2$ and μ is an arbitrary scale parameter introduced with zeta regularization. We compute the functions f_0 and f_1 with some range of the deformation parameter ρ . It is interesting to note that in $R^4 \times \tilde{S}^2$ the divergent part in (1) disappears with $\rho \simeq 0.415$ yielding the finite value $E \simeq 1.47 \cdot 10^{-5}$. (Here and throughout the paper we set $r = 1$). We compare our results with those obtained in flat spaces for a scalar field satisfying in two dimensions to the Dirichlet, von Neumann and periodic boundary conditions.

Let us begin with the scalar Laplacian eigenvalues on a deformed sphere $\tilde{S}^2$. The metric on $\tilde{S}^2$ can be expressed in the form

$$ds^2 = dx_0^2 + \sin^2 x_0 \rho^{-2} d\psi^2, \quad 0 \leq \psi \leq 2\pi$$

where ρ is a positive constant parameter. This space can be also obtained by identification of points with polar angles α and $\alpha + 2\pi/\rho$ on round S^2 with a maximally symmetric metric. The $\tilde{S}^2$ has two singular points $x_0 = 0$ and $x_0 = \pi$. Let us perform the Fourier decomposition with respect to the coordinate ψ .

$$\phi(x_0, x_i) = \sum_{l=-\infty}^{\infty} f_{(l)}(x_0) \exp(il\psi) \quad (2)$$

Substituting the decomposition (2) in the eigenvalue equation

$$\Delta\phi = -\lambda\phi$$

we obtain the following ordinary differential equation

$$[\partial_0^2 + \text{ctg} x_0 \partial_0 - \frac{\rho^2 l^2}{\sin^2 x_0}] f_{(l)} = -\lambda f_{(l)}. \quad (3)$$

Equations of this type were considered in our previous papers [14]. We shall drop the subscripts (l) for a while. Let us change the independent and dependent variables

$$f = h \sin^{|l|\rho}(x_0), \quad z = \frac{1}{2}(\cos x_0 + 1).$$

The $|l|$ denotes absolute value of l . The equation (3) then takes the form

$$z(z-1)h'' + 2(\rho|l|+1)(z-\frac{1}{2})h' + (\rho|l|(\rho|l|+1)-\lambda)h = 0. \quad (4)$$

Prime denotes differentiation with respect to z . According to the general prescription [16] let us express h as the power series

$$h(z) = \sum_{k=0} \alpha_k z^k. \quad (5)$$

Substitution of (5) in (4) gives a recurrent condition on the coefficients α_k

$$\alpha_{k+1} = \alpha_k \frac{k(k-1) + 2(\rho|l|+1)k + \rho|l|(\rho|l|+1) - \lambda}{(k+1)(k+\rho|l|+1)}. \quad (6)$$

The denominator of (6) is positive for all k . The eigenfunctions h_k can be found by imposing the condition on the numerator of (6) to be equal to zero. We obtain the eigenvalues

$$\lambda_{l,k} = (k + \rho|l| + \frac{1}{2})^2 - \frac{1}{4} \quad (7)$$

where we restored the dependence on the index l . (Eq. 7 has been also considered in Ref. [15]).

In the case of the unit round sphere S^2 with $\rho = 1$ we obtain from (7)

$$\lambda_{(l)k} = -(k + |l|)(k + |l| + d) = -n(n + d), \quad n = k + |l|$$

Thus equation (6) reproduces the correct eigenvalues of the scalar Laplace operator on the unit round S^2 . One can also verify that the degeneracies have the correct values.

With zeta regularization the Cazimir energy density for a scalar field on $R^{m+1} \times \tilde{S}^2$ can be written as

$$E = \lim_{s \rightarrow -1} \frac{\Gamma((s-m)/2) \mu^{2\epsilon}}{2(4\pi)^{m/2} \Gamma(s/2)} \left(2 \sum_{l=1}^{\infty} \sum_{k=0}^{\infty} \lambda_{l,k}^{-s/2} + \sum_{k=0}^{\infty} \lambda_{0,k}^{-s/2} \right) \quad (8)$$

where $\epsilon = (1+s)/2$. Making use of the Hermite formula [16]

$$\zeta(z, q) = \frac{q^{-z}}{2} + \frac{q^{1-z}}{z-1} + 2 \int_0^{\infty} dx \sin(z \arctan(x/q)) \frac{(q^2 + x^2)^{-z/2}}{e^{2\pi x} - 1}$$

and taking the expansion

$$\sum_{n=0}^{\infty} (n+q)^l ((n+q)^2 + M)^{\frac{m+s}{2}} = \sum_{k=0}^{\infty} \frac{(-1)^k \Gamma(k - (m+s)/2)}{\Gamma(-(m+s)/2) k!} \zeta(2k - l - m - s, q) M^k$$

we rewrite (8) as

$$E = \lim_{s \rightarrow -1} \sum_{r=0}^{\infty} \frac{\Gamma(r + (s-m)/2) \mu^{2\epsilon}}{2(4\pi)^{m/2} \Gamma(s/2) 4^r r!} \left(P(s-m+2r, \rho) + F(s-m+2r, \rho) \right) \quad (9)$$

where

$$\begin{aligned}
P(z, \rho) &= \rho^{-z} \zeta(z, 1 + 1/(2\rho)) + \frac{2}{z-1} \rho^{1-z} \zeta(z-1, 1 + 1/(2\rho)) + \zeta(z, 3/2), \\
F(z, \rho) &= 2 \sum_{p=0}^{\infty} (-1)^{p+1} c_p(z) \sum_{n=0}^{\infty} \frac{\Gamma(n + z/2)}{\Gamma(z/2) n!} \rho^{-2p-2n-z-1} \\
&\quad \times \zeta(2p + 2n + z + 1, 1 + 1/(2\rho)) \zeta(-2p - 2n - 1).
\end{aligned} \tag{10}$$

The coefficients c_p are determined from

$$\sin(z \arctan(x)) = \sum_{p=0}^{\infty} c_p(z) x^{2p+1}$$

The expression (9) involves the divergent terms coming from the poles of $\Gamma(z)$ at $z = -k$ ($k = 0, 1, 2, \dots$)

$$\Gamma(-k + \epsilon) = \frac{(-1)^k}{k!} \left(\frac{1}{\epsilon} + \psi(k+1) + O(\epsilon) \right) \tag{11}$$

and the pole of $\zeta(z, q)$ at $z = 1$

$$\zeta(1 + 2\epsilon, q) = \frac{1}{2\epsilon} - \psi(q) + O(\epsilon). \tag{12}$$

However, in odd dimensions the divergent term $1/(z-1)$ at $z = 1$ cancels the pole of zeta function yielding the finite Casimir energy. We find, omitting the details, from (9)-(12) the following numerical values for the Casimir energy E_m .

	$\rho = 0.4$	$\rho = 0.8$	$\rho = 1$	$\rho = 2$
E_0	0.07385	- 0.09987	- 0.13246	- 0.1990,
E_2	$16.1 \cdot 10^{-4}$	$- 1.53 \cdot 10^{-4}$	$- 4.78 \cdot 10^{-4}$	$- 11.4 \cdot 10^{-4}$. (13)

In even dimensions, for E_m given in the form (1) we compute both $f_0(m, \rho)$ and $f_1(m, \rho)$.

For $m = 1$

	$\rho = 0.4$	$\rho = 0.8$	$\rho = 1$	$\rho = 2$
$f_0(1, \rho)$	- 0.00144	- 0.00216	- 0.00265	- 0.0063,
$f_1(1, \rho)$	- 0.0023	- 0.0097	- 0.0117	- 0.0179. (14)

For $m = 3$

$f_0(3, \rho)$	$1.2 \cdot 10^{-6}$	$- 2.56 \cdot 10^{-5}$	$- 4.02 \cdot 10^{-5}$	$- 21.8 \cdot 10^{-5},$
----------------	---------------------	------------------------	------------------------	-------------------------

$$f_1(3, \rho) \quad 2.81 \cdot 10^{-5} \quad -6.18 \cdot 10^{-5} \quad -9.15 \cdot 10^{-5} \quad -18.1 \cdot 10^{-5}. \quad (15)$$

Now some notes are in order. To extract the effect of deformation we should redefine r such that the volume on $\tilde{S}^2$ is the same as that of S^2 with a unit radius. In this case, we have to consider the quantity $\bar{E}_m = E/\rho^{m+1}$. It is easy to see from (13) that $\bar{E}_0$ and $\bar{E}_2$ take the minimal values at $\rho = 1$. Thus, sphere S^2 tends to keep its shape at least for the slight deformation. A detailed study shows that this is the only minimum of $\bar{E}_m$. The asymptotic behaviours of $\bar{E}_m$ ($m = 0, 2$) are

$$\bar{E}_m \sim \rho^{-m-2}, \quad \rho \rightarrow 0,$$

$$\bar{E}_m \sim -\rho^{-m-1} \log \rho \quad \rho \rightarrow \infty.$$

Indeed, the singularity at $\rho = 0$ gives rise to the repulsive force, tending to expand $\tilde{S}^2$. Since $\bar{E}_1$ and $\bar{E}_3$ have the divergences and depend on an arbitrary scale μ , some renormalization procedure should be used to obtain the finite result. However, one can see from (15) that $f_0(3, \rho)$ becomes equal to zero with the certain value of $\rho = \rho_0$ in the region $0.4 < \rho < 0.6$ and the divergent part is removed from E_3 . This reflects the fact that the conformal anomaly in $R^4 \times \tilde{S}^2$ is canceled with the deformation parameter ρ_0 .

Finally, we compare the results in (13) with those obtained for a scalar massless field in general three- and five-dimensional hyper-cuboidal regions with two sides of finite length a_1, a_2 and one and three sides with length $L \gg a_1, a_2$ respectively. In the case of periodic boundary conditions the vacuum energy as a function of a_2/a_1 has the minimal value when $a_2 = a_1$ and is always negative. The energy obtained with von Neumann boundary conditions [2] also exhibits this behaviour. With Dirichlet boundary conditions the energy takes the positive maximal values at $a_2/a_1 = 1$ for both three- and five-dimensional spaces and tends to negative infinity when $a_2/a_1 \gg 1$ [2]. Thus, we can see that in all these models the behaviour of the vacuum energy is quite different from that of the energy in $R^1 \times \tilde{S}^2$ and $R^3 \times \tilde{S}^2$ spaces.

Acknowledgements

This work was partially supported by the Russian Foundation for Fundamental Studies, grant 93-02-14378. One of us (DV) is grateful to Ignati Grigentch for colaboration on Laplace operator on deformed spheres and to Sergio Zerbini, Guido Cognola and Luciano Vanzo for hospitality at Trento and discussions. We are also grateful to D. Fursaev for pointing out Ref. [15].

References

- [1] H.B.G. Casimir, J.Chim. Phys. **46** (1949) 407;
R. Balian and B. Duplaintier, Ann. Phys. **112** (1978) 165;
K.A.Milton, L.L. DeRaad Jr. and J. Schwinger, Ann. Phys. **115** (1978) 388;
G. Kennedy, J. Phys. A **11** (1978) L173;
D. Deutsch and P. Candelas, Phys. Rev. D**20** (1979) 3063.
- [2] J. Ambjorn and S. Wolfram, Ann. Phys. **147** (1983) 1;
J. Ambjorn and S. Wolfram, Ann. Phys. **147** (1983) 33.
- [3] S. Sen, J. Math. Phys. **22** (1981) 2968;
S. Sen, J. Math. Phys. **25** (1984) 2000.
- [4] E. Elizalde, Phys. Lett. **213 B** (1988) 477;
E. Elizalde and A. Romeo, J. Math. Phys. **33** (1989) 1133;
E. Elizalde, Z. Phys. C**44** (1989) 471.
- [5] J. S. Dowker and G. Kennedy, J. Phys. A **11** (1978) 895;
G. Kennedy, R. Critchly and J. S. Dowker, Ann. Phys. **125** (1980) 346;
S. Tadaki and S. Takagi, Prog. Theor. Phys. **75** (1986) 262;
S. A. Gundersen and F. Ravndal, Ann. Phys. **182** (1988) 90;
K. Kirsten, J. Phys. A **24** (1991) 3281 .
- [6] S. K. Blau, M. Visser and A. Wipf, Nucl. Phys. **B310** (1988) 163.
- [7] G. Plunien, B.Muller and W.Greiner, Phys.Rep.**134** (1986) 87.
- [8] P. Candelas and S. Weinberg, Nucl. Phys. **B237** (1984) 397;
M.A. Awada and D.J. Toms, Nucl. Phys. **B245** (1984) 161;
D. Birmingham and S. Sen, Ann. Phys. **172** (1986) 451;
T.C. Shen and J. Sobczyk , Phys. Rev. D**36** (1987) 397.
- [9] J. S. Dowker, Commun. Math. Phys. **162** (1994) 633;
J. S. Dowker, J. Math. Phys. **35** (1994) 4889.
- [10] K. Schleich and D.H. Witt, Nucl. Phys. **B402** (1993) 411.
- [11] J.S.Dowker, J. Phys. **A10** (1977) 115;
J.S.Dowker, Phys. Rev. **D36** (1987) 3742;
A.Vilenkin, Phys. Rev. **D23** (1981) 852;

- S.Deser, R.Jackiw and G.t'Hooft, Ann. Phys. **152** (1984) 220;
P. de Sousa Gerbert and R.Jackiw, Commun. Math. Phys. **124** (1989) 229;
V.D.Skarzhinski, D.D.Harary and U.Jasper, Phys. Rev. **D49** (1994) 755.
- [12] G.Cognola, K.Kirsten and L.Vanzo, Phys. Rev. **D49** (1994) 1027;
G.Cognola and L.Vanzo, J. Math. Phys. **35** (1994) 3109;
D.V.Fursaev and G.Miele, Phys. Rev. **D49** (1994) 987. D.V.Fursaev, Phys. Lett. **334 B** (1994) 53.
- [13] J.Cheeger, J. Diff. Geom. **18** (1983) 575;
J.S.Dowker, J. Math. Phys. **30** (1989) 770;
J.S.Dowker, Class. Quantum Grav. **11** (1994) 557;
P.Chang and J.S.Dowker, Nucl. Phys. **B395** (1993) 407.
- [14] N.Shtykov and D.V.Vassilevich, TIT preprint TIT/HEP-265/94;
I.P.Grigentch and D.V.Vassilevich, Nuovo Cimento **107A** (1994) 227.
- [15] D. Fursaev, Class. Quantum Grav. **11** (1994) 1431.
- [16] E.T.Whittaker and Watson , A course of modern analysis, Cambridge University Press, 1927.